



FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2014/2015

Ana Vera Cunha Bessa da Costa
Severity of disease scoring systems and
mortality after non-cardiac surgery

março, 2015

FMUP



FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

Ana Vera Cunha Bessa da Costa
Severity of disease scoring systems and
mortality after non-cardiac surgery

Mestrado Integrado em Medicina

Área: Anestesiologia

Tipologia: Dissertação

Trabalho efetuado sob a Orientação de:
Doutor Fernando José Pereira Alves Abelha

Trabalho organizado de acordo com as normas da revista:
Journal of critical care

março, 2015

FMUP

Eu, Ana Vera Cunha Bessa da Costa, abaixo assinado, nº mecanográfico 090801219, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 23/02/2015

Assinatura conforme cartão de identificação:

Ana Vera Cunha Bessa da Costa

Projecto de Opção do 6º ano – DECLARAÇÃO DE REPRODUÇÃO

NOME

Ana Vera Cunha Bessa da Costa

CARTÃO DE CIDADÃO OU PASSAPORTE (se estrangeiro)

E-MAIL

TELEFONE OU TELEMÓVEL

13967049

mimed09231@med.up.pt

918130002

NÚMERO DE ESTUDANTE

DATA DE CONCLUSÃO

090801219

23 de março de 2015

DESIGNAÇÃO DA ÁREA DO PROJECTO

Anestesiologia

TÍTULO DISSERTAÇÃO

Severity of disease scoring systems and mortality after non-cardiac surgery

ORIENTADOR

Prof. Doutor Fernando José Abelha

É autorizada a reprodução integral desta Dissertação para efeitos de investigação e de divulgação pedagógica, em programas e projectos coordenados pela FMUP.

Faculdade de Medicina da Universidade do Porto, 23/03/2015

Assinatura conforme cartão de identificação: Ana Vera Cunha Bessa da Costa

À Graça e ao Manuel

Severity of disease scoring systems and mortality after non-cardiac surgery

Ana Vera Costa¹, Pedro Videira Reis², Gabriela Sousa², Ana Martins Lopes², Fernando José Abelha^{2,3}

¹ Faculty of Medicine, University of Porto, Alameda Professor Hernâni Monteiro, Hospital de S. João, 4200-319 Porto, Portugal

²Department of Anesthesiology, Centro Hospitalar São João, Alameda Professor Hernâni Monteiro, Hospital de S. João, 4200-319 Porto, Portugal

³Department of Anesthesiology and Perioperative Medicine - University of Porto Faculty of Medicine, Alameda Professor Hernâni Monteiro, Hospital de S. João, 4200-319 Porto, Portugal

E-mail addresses:

Ana Vera Costa: mimed09231@med.up.pt

Pedro Videira Reis: pedrojsreis@hotmail.com

Gabriela Sousa: gscsousa@gmail.com

Ana Martins Lopes: analopes13@gmail.com

Corresponding author:

Fernando José Abelha

Telephone: 00351917304213

E-mail: fernando.abelha@gmail.com

Fax number: 00351967599051

Postal address: Alameda da República, 731, Gulpilhares – 4405-696 Vila Nova de Gaia, Portugal

Abstract

Purpose: There are still high death rates related to surgeries mainly linked to postoperative period. Mortality prediction is one of the greatest challenges posed in intensive care and severity of disease scoring systems are the main tool available. The aim of this study was to evaluate predictors for postoperative mortality.

Methods: all adult patients who underwent non-cardiac surgery, admitted in a surgical intensive care unit (SICU) for longer than 12 hours between January 2006 and July 2013 were evaluated. Univariate analysis was carried using Mann-Whitney test, Chi-Square or Fischer's exact test. Multivariate analysis with logistic regression was performed with calculation of Odds Ratio (OR) with 95% confidence interval (CI).

Results: 4398 patients met the inclusion criteria. Mortality in SICU was 1.4%. Patients dying in the SICU had higher score in severity of disease scoring systems and length of stay. Multivariate analyses find 5 independent predictors of mortality in the SICU: APACHE II (OR 1.24, CI 1.17-1.30), emergent surgery (OR 4.10, CI 2.12-7.94), postoperative serum sodium (OR 1.06, CI 1.01-1.11), serum bicarbonate (OR 0.89, CI 0.82-0.96), and FiO₂ (OR 14.31, CI 3.01-68.07).

Conclusion: independent predictors of SICU's mortality were APACHE II, emergent surgery, postoperative serum bicarbonate, serum sodium and FiO₂.

Keywords: postoperative mortality, severity of disease scoring systems, APACHE II
SAPS II, surgical intensive care unit, non-cardiac surgery

Abbreviations

ASA-PS: American Society of Anesthesiologists physical status; CI: confidence interval; GCS: glasgow coma scale; HR: heart rate; ICU: intensive care unit; LOS: length of stay; MAP: mean arterial pressure; OR: odds ratio; PACU: Post-Anesthesia Care Unit; RCRI: Revised Cardiac Risk Index; RR: respiratory rate; SAPS: Simplified Acute Physiology Score; SICU: surgical intensive care unit; SP: systolic pressure

Conflict of interest: none

Submission declaration: the work described will be published as an academic thesis, Faculty of Medicine of Porto, Portugal, 2015. Preliminary data for this study was presented at the congress of the Portuguese Society of Anesthesiology, Cascais, Portugal (13 march 2015).

Introduction

It is estimated that every year 234.2 million people will be submitted to surgery worldwide (1).

According to the 2012 European Surgical Outcomes Study (EuSOS), 4% of patients died before hospital discharge (2) and postoperative one-year mortality was 5.5% (3).

The majority of deaths occur in older patients who undergo major emergent surgery and who have severe coexisting diseases as well in those patients that develop complications (4-7).

There are several risk factors described for morbidity and mortality after surgery, which may be divided into three categories: patient-related, surgery-related and anesthesia-related. The risks of surgery and anesthesia are low for the majority of patients and evidence increasingly suggests that postoperative complications have a major impact on mortality (4, 5, 8). Developed countries have major morbidity due to postoperative complications (12 % in United States) (9). Thereof with aging and consequent increase of the patient's co-morbidities, as well as with increasing numbers of patients, there is an increase in postoperative morbidity and mortality (5, 10).

Half of the postoperative adverse events were identified as avoidable (11). Reducing rates of postoperative complications and their effective management may be one approach in reduce postoperative mortality (4, 5, 9). Immediate postoperative care allows a close monitoring and early intervention in order to prevent early postoperative complications and deaths. Seriously ill or at increased risk patients' may require more extensive monitoring in a surgical intensive care unit (SICU) which may contribute to a better outcome, decreasing morbidity and mortality. However there are few SICU's beds and high costs of their use (12, 13).

The study of outcome in critical care patients has been primarily focused on hospital survival and the resource utilization adjusted according to the severity of illness.

Intensive care unit (ICU) case fatality rates strongly depend on the severity of illness of the individual in the patient population being analyzed (14).

To improve postoperative care, severity of disease scoring systems were used in order to predict the prognosis and estimate the morbidity and mortality of patients. Acting as outcome predictors they allow comparisons and the calculation of the expected mortality. Acute Physiology and Chronic Health Evaluation II (APACHE II) and Simplified Acute Physiology Score II (SAPS II) are two worldwide used severity of disease scoring systems (15-17).

The severity of scoring systems may be used to predict mortality with the calculation of the standardized mortality ratio (SMR), the ratio of observed to predicted mortality, which is used as an indicator of the quality of intensive care (18-20), although some authors argue that the SMR should not be used to measure the quality of care (21, 22). Several risk indices have been developed over the past years based on the relationship between comorbidities and perioperative morbidity and mortality. The Revised Cardiac Risk Index (RCRI) have become well known and although it isn't a severity of disease score has been used to predict an increased risk of cardiac complications after surgery being incorporated in guidelines for the study of preoperative risk factors (23, 24). The aim of the present study was to evaluate the determinants of mortality using parameters included in severity of disease scoring systems in a cohort of critical surgical patients.

Methods

Data collection

The full study protocol was approved by the research ethics committee of the hospital. This prospective cohort study was carried out in the multidisciplinary Post-Anesthesia Care Unit (PACU) at the Hospital São João, an 1124-bed community teaching hospital in Porto, Portugal. Included in the PACU was a Surgical Intensive Care Unit (SICU) with five beds to which critically ill surgical patients are admitted and are closely monitored and treated.

All patients admitted at the SICU who underwent elective or emergent non-cardiac surgery between 1st January 2006 and 19th July 2013 were eligible for inclusion. Patients less than 18 years old, medical patients, re-admittance for the same medical reason during the studied period and SICU length of stay (LOS) lower than 12 hours were excluded.

The following variables were recorded at admission to the SICU: age, type of admission (elective or non-elective surgery) and mechanical ventilation.

For all patients we recorded the SICU LOS and the mortality in the SICU. Major cardiac event was defined as at least one of the following: acute myocardial infarction, acute pulmonary edema, primary cardiac arrest, ventricular fibrillation/other ventricular arrhythmias, complete heart block were assessed (25).

Organ insufficiency (considering presence of at least one organ failure defined by APACHE II) and renal insufficiency (considering creatinine > 2 mg/dL and/or oliguria of <500 ml / day) was also evaluated.

APACHE II and SAPS II were calculated. All variables and parameters included in APACHE II and SAPS II scores were evaluated separately (15, 17).

RCRI was also evaluated using criteria developed by Lee et al: high-risk surgery (intraperitoneal, intrathoracic, or suprainguinal vascular procedures), history of ischemic heart disease, history of congestive heart disease, preoperative insulin therapy, preoperative serum creatinine >2.0 mg/dL and history of cerebrovascular disease (23).

Statistical analysis

Descriptive analysis of variables was used to summarize data.

Kolmogorov-Smirnov Test for normality of the underlying population was performed.

The Mann-Whitney U test, the Chi-square and Fisher's exact test were used in the univariate analyses to compare continuous variables and proportions, respectively.

To assess independent predictive factors of postoperative mortality, multivariate analysis were performed with multiple binary logistic regression. After applying the Bonferroni's correction for multiple comparisons, all the variables included in severity of disease scoring systems that had $p \leq 0.001$ in the univariate analyses were entered in a logistic multiple regression binary analysis with forward elimination method to examine covariate effects on mortality calculating an odds ratio (OR) and 95% confidence interval (CI)

The statistical software SPSS version 22.0 for Windows (SPSS, Chicago, IL) was used to analyze the data.

Results

During the study period there were 4561 admissions in the SICU and 4398 patients met the inclusion criteria. A total of 163 patients were excluded: 53 had a length of SICU stay lower than 12 hours, 42 were admitted more than once, 38 were younger than 18 years old and 30 were admitted for medical reasons. Sixty patients (1.4%) died in the SICU.

Table 1 display the characteristics of all patients enrolled in the study and the comparison between patients who survive and patients that have died during SICU stay.

The median age was 65 years (IQR 54 -74 years) and 61% were male.

Thirteen percent of patients were admitted after non-elective surgery.

The median postoperative length of stay for all patients was 20 hours (IQR 16-42 hours).

One hundred and seven patients (2.4%) developed major cardiac event that include at least one of the following events: acute myocardial infarction (n=59), acute pulmonary edema (n=34), primary cardiac arrest (n=19), ventricular fibrillation/other ventricular arrhythmias (n=8) and complete heart block (n=4).

In univariate analyses patients that died were older (median 73 vs 65, $p<0.001$), were more likely to have been submitted to an emergent surgery (60% vs 12%, $p<0.001$), had more frequently mechanical ventilation on admission (83% vs 30%, $p<0.001$), renal insufficiency (35% vs 6%, $p<0.001$) and organ insufficiency (40% vs 15%, $p<0.001$).

Patients that have died in SICU had lower hematocrit (median 29 vs 33, $p<0.001$), lower body temperature (median 34.0 vs 35.8, $p<0.001$), lower systolic pressure (median 77 vs 122, $p<0.001$), lower mean arterial pressure (median 54 vs 85, $p<0.001$), higher heart rate (median 112 vs 83, $p<0.001$), higher respiratory rate (median 16 vs 14, $p<0.001$), higher urea serum concentration (median 45 vs 30, $p<0.001$), higher serum creatinine (median 15.6 vs 8.2, $p<0.001$), higher total bilirubin (median 6 vs 4, $p<0.001$), higher FiO₂ (median 0.52 vs 0.40, $p<0.001$), lower PaO₂ (median 98 vs 100, $p=0.039$), higher PaCO₂ (median 42.7 vs 39.4, $p=0.001$), lower serum bicarbonate (median 19.4 vs 22.0, $p<0.001$), lower pH (median 7.28 vs 7.40, $p<0.001$), higher serum sodium (median 145 vs 140, $p<0.001$) and had more frequently a glasgow coma scale < 9 (8% vs 1%, $p<0.001$).

Table 2 displays severity of disease scores, length of stay and major cardiac events.

Patients that died had higher scores of APACHE II (median 22 vs 8, $p<0.001$) and SAPS II (median 44 vs 18, $p<0.001$) and had more frequently major cardiac events

(28% vs 2%, $p<0.001$) and had a longer stay (median 46 vs 20, $p<0.001$) compared to survival group.

As displayed in table 3, patients not surviving had more frequently RCRI scores of more than 2 (18% versus 7%, $p=0.004$). These patients were submitted more frequently to a high-risk surgery (80% vs 54%, $p<0.001$).

In table 4 are the results of the multivariate analyses that showed that APACHE II (OR 1.24, 95% CI 1.17-1.30), type of admission as emergent (OR 4.10, 95% CI 2.12-7.94), FiO₂ (OR 14.31, 95% CI 3.01-68.07) and serum sodium (OR 1.06, 95% CI 1.01-1.11) are independent predictors of mortality while serum bicarbonate (for each 1 mmol/L increase) was associated with reduced risk of death (OR 0.89, 95% CI 0.82-0.96).

Discussion

In the present study, we assessed the mortality rate of a cohort of surgical critical patients after major surgery and studied predictors of mortality analyzing all variables included in the severity of score disease APACHE II and SAPS II.

The principal findings of this study were as follows: that the mortality rate in the SICU was 1.4%; that patients not surviving after SICU admission had higher severity of disease scores according to SAPS II and APACHE II, they stayed longer in the SICU and had more frequently major cardiac events; almost all variables included in the severity of disease scoring systems were different in the group of patients not surviving in SICU. When adjusted for each other and maintaining severity of disease scores in the equation only APACHE II, emergent surgery type of admission, inspired fraction of oxygen and serum sodium were considered independent factors for mortality while a higher serum bicarbonate was associated with a reduction of mortality.

Early identification of patients at a higher risk of mortality may be important to promote treatment and orientation strategies that may preclude case fatalities (4, 6). Severity of disease scoring systems may allow us to estimate mortality risk in surgical critical patients (15-17) and identifying a group of patients at increased risk of dying.

Previous studies have been focused on identifying predictors of postoperative mortality and morbidity evaluating and quantifying comorbidities, perioperative factors and the presence of postoperative complications (3-6, 25-33), but no studies have attempted to identify predictors within routine physiological and analytical postoperative parameters included in severity of disease scoring systems.

Type of surgery is a variable that has been studied and clearly found to be related to mortality (6, 7, 10, 30, 34) and our results indicate as well that emergent surgery patients had a higher mortality rate. It seems that patients undergoing non-elective surgeries are likely to have a worse prognosis since they were more severely ill with a higher grade of comorbidities with a higher probability of having a less functional reserve. Because of that these patients may be submitted to more complex surgeries and they usually require a careful intraoperative care (10, 35). In our study non-elective emergent surgery was considered an independent predictor of mortality, increasing the probability of dying in the SICU almost 4 times.

In multivariate analysis FIO₂ was another independent predictor of mortality. Higher FIO₂ is frequently required in patients with impaired tissue oxygenation trying to avoid the harmful effects of hypoxia. In fact it is well documented that the PaO₂/FIO₂ ratio is

associated with mortality (36, 37). In our study we did not studied this equation fraction but the isolated FiO₂ parameter may be considered as a relevant surrogate indicator of that fraction. In a previous study higher FiO₂ remained an independent predictor of mortality even after adjustment for PaO₂/FiO₂ ratio (38), suggesting poor prognosis not only because these patients are more severely ill with impaired tissue oxygenation, but also because of hyperoxia and ventilation side-effects (38-40).

Others have found serum sodium to be a reliable risk factor for mortality (41-45) and it is not surprisingly that in this study serum sodium was considered an independent risk factor for mortality. Hyponatremia is a common complication in critically ill patients such they may be unconscious, intubated or sedated and may invariably denotes hyperosmolar state and transiently intracellular dehydration (46).

The multivariate analysis of independent variables showed that higher serum bicarbonate was associated with a reduction of mortality for each mmol increment having a positive impact on survival. This result may indicate that a lower serum bicarbonate levels was associated with metabolic acidosis and consequently with case fatalities that has been shown by others (47-50). Although the deleterious impact of lower serum bicarbonate is well known, both lower and higher serum bicarbonates may be associated with increased all-cause mortality as a result of the well documented consequences of acid-base abnormalities that have been associated with adverse outcomes such as increased mortality (51). However, a recent retrospective analysis shows that acidosis itself had no relation with poor outcome which was more dependent on severe conditions associated with acidosis (52).

Age have also been found to have a clear association with postoperative mortality (3, 6, 53) and that is why is not surprisingly that in our study patients that died in the SICU were older.

In order to stratify the preoperative risk of patients we relied on the RCRI. Patients that died had more frequently a score greater or equal than 2 but only high-risk surgery were considered a risk factor for mortality and it is interesting that comorbidities included in RCRI did not appear to be related to mortality. Perhaps in this particular group of major surgery patients the burden of surgery was more relevant than comorbidities.

Patients with prolonged SICU LOS had higher mortality as LOS is needed in patients who develop postoperative complications and those who were more severely ill (7, 25-27).

Data suggest that complication rate varies between 7% and 11% (54). The occurrence of postoperative cardiac events is not uncommon, occurring in 2.4% in this study.

Furthermore mortality group had significantly more major cardiac events, corroborating previous studies showing that patients with postoperative major cardiac events had higher mortality (25, 26). Indeed previous studies show that cardiovascular complications are the leading causes of morbidity in older patients (10) and may be associated with 42% of deaths (34).

We know that many other complications are related to increased risk of mortality (4, 9, 55) and that many of the surviving patients suffer major complications with a variety of morbidity and the consequent reduction of the health and quality of life (26, 56).

Study limitations

Besides the limitations inherent to a retrospective cohort study, others are present on the design of this study. Preoperative risk assessment is roughly based on three broad but connected categories including several risk factors: surgery-related, patient-related or dependent on patient's functional status. Not knowing the pre-existing conditions of patients beyond the comorbidities present in the Revised Cardiac Risk Index probably may also limit the value of conclusions, because comorbidities others than those we have studied may be closely related and may have a clear influence on physiological parameters included in the severity of disease scoring systems (for example chronic obstructive pulmonary disease and PaO₂). The lack of an American Society of Anesthesiologists physical status (ASA-PS) for our sample population was also questionable. Risk prediction models for intraoperative and postoperative mortality have included the ASA-PS classification as one strong predictor (24, 57-59).

Other limitation of the study is the missing data about in-hospital mortality what may be viewed as an important limitation since severity of disease scores and its standard mortality rates are related to hospital mortality not to ICU mortality.

Furthermore, neither intraoperative hemodynamic parameters nor other postoperative complications beyond organ insufficiency, renal insufficiency and major cardiac events were evaluated in our study and this may be viewed as a limitation that certainly had influenced outcome and mortality.

Conclusions

In conclusion postoperative mortality during SICU stay was 1.4%. Fatalities cases had significantly higher scores in severity of disease scoring systems as well as had a longer SICU stay and more major cardiac events. We have identified independent risk factors for mortality: APACHE II, FiO₂, serum sodium, type of admission while higher serum bicarbonate was associated with a reduction of mortality.

References

1. Weiser TG, Regenbogen SE, Thompson KD, Haynes AB, Lipsitz SR, Berry WR, et al. An estimation of the global volume of surgery: a modelling strategy based on available data. *Lancet*. 2008;372(9633):139-44.
2. Pearse RM, Moreno RP, Bauer P, Pelosi P, Metnitz P, Spies C, et al. Mortality after surgery in Europe: a 7 day cohort study. *Lancet*. 2012;380(9847):1059-65.
3. Monk TG, Saini V, Weldon BC, Sigl JC. Anesthetic management and one-year mortality after noncardiac surgery. *Anesthesia and analgesia*. 2005;100(1):4-10.
4. Khuri SF, Henderson WG, DePalma RG, Mosca C, Healey NA, Kumbhani DJ, et al. Determinants of long-term survival after major surgery and the adverse effect of postoperative complications. *Annals of surgery*. 2005;242(3):326-41.
5. Ghaferi AA, Birkmeyer JD, Dimick JB. Variation in hospital mortality associated with inpatient surgery. *The New England journal of medicine*. 2009;361(14):1368-75.
6. Pearse RM, Harrison DA, James P, Watson D, Hinds C, Rhodes A, et al. Identification and characterisation of the high-risk surgical population in the United Kingdom. *Critical care*. 2006;10(3):R81.
7. Abelha FJ, Castro MA, Landeiro NM, Neves AM, Santos CC. Mortality and length of stay in a surgical intensive care unit. *Revista brasileira de anestesiologia*. 2006;56(1):34-45.
8. Arbous MS, Grobbee DE, van Kleef JW, de Lange JJ, Spoormans HH, Touw P, et al. Mortality associated with anaesthesia: a qualitative analysis to identify risk factors. *Anaesthesia*. 2001;56(12):1141-53.
9. Hall BL, Hamilton BH, Richards K, Bilimoria KY, Cohen ME, Ko CY. Does surgical quality improve in the American College of Surgeons National Surgical Quality Improvement Program: an evaluation of all participating hospitals. *Annals of surgery*. 2009;250(3):363-76.
10. Leung JM, Dzankic S. Relative importance of preoperative health status versus intraoperative factors in predicting postoperative adverse outcomes in geriatric surgical patients. *Journal of the American Geriatrics Society*. 2001;49(8):1080-5.
11. Kable AK, Gibberd RW, Spigelman AD. Adverse events in surgical patients in Australia. *International journal for quality in health care : journal of the International Society for Quality in Health Care / ISQua*. 2002;14(4):269-76.

12. Weissman C. The enhanced postoperative care system. *Journal of clinical anesthesia*. 2005;17(4):314-22.
13. Simpson JC, Moonesinghe SR. Introduction to the postanesthetic care unit. *Perioperative medicine*. 2013;2(1):5.
14. Halpern NA, Pastores SM, Greenstein RJ. Critical care medicine in the United States 1985-2000: an analysis of bed numbers, use, and costs. *Critical care medicine*. 2004;32(6):1254-9.
15. Knaus WA, Draper EA, Wagner DP, Zimmerman JE. APACHE II: a severity of disease classification system. *Critical care medicine*. 1985;13(10):818-29.
16. Vincent JL, Moreno R. Clinical review: scoring systems in the critically ill. *Critical care*. 2010;14(2):207.
17. Le Gall JR, Lemeshow S, Saulnier F. A new Simplified Acute Physiology Score (SAPS II) based on a European/North American multicenter study. *Jama*. 1993;270(24):2957-63.
18. Sirio CA, Shepardson LB, Rotondi AJ, Cooper GS, Angus DC, Harper DL, et al. Community-wide assessment of intensive care outcomes using a physiologically based prognostic measure: implications for critical care delivery from Cleveland Health Quality Choice. *Chest*. 1999;115(3):793-801.
19. Jarman B, Pieter D, van der Veen AA, Kool RB, Aylin P, Bottle A, et al. The hospital standardised mortality ratio: a powerful tool for Dutch hospitals to assess their quality of care? *Quality & safety in health care*. 2010;19(1):9-13.
20. Breslow MJ, Badawi O. Severity scoring in the critically ill: part 2: maximizing value from outcome prediction scoring systems. *Chest*. 2012;141(2):518-27.
21. Lilford R, Pronovost P. Using hospital mortality rates to judge hospital performance: a bad idea that just won't go away. *Bmj*. 2010;340:c2016.
22. Mohammed MA, Deeks JJ, Girling A, Rudge G, Carmalt M, Stevens AJ, et al. Evidence of methodological bias in hospital standardised mortality ratios: retrospective database study of English hospitals. *Bmj*. 2009;338:b780.
23. Lee TH, Marcantonio ER, Mangione CM, Thomas EJ, Polanczyk CA, Cook EF, et al. Derivation and prospective validation of a simple index for prediction of cardiac risk of major noncardiac surgery. *Circulation*. 1999;100(10):1043-9.
24. Kristensen SD, Knuuti J, Saraste A, Anker S, Botker HE, De Hert S, et al. 2014 ESC/ESA Guidelines on non-cardiac surgery: cardiovascular assessment and management: The Joint Task Force on non-cardiac surgery: cardiovascular assessment

- and management of the European Society of Cardiology (ESC) and the European Society of Anaesthesiology (ESA). *European journal of anaesthesiology*. 2014.
25. Maia PC, Abelha FJ. Predictors of major postoperative cardiac complications in a surgical ICU. *Revista portuguesa de cardiologia : orgao oficial da Sociedade Portuguesa de Cardiologia = Portuguese journal of cardiology : an official journal of the Portuguese Society of Cardiology*. 2008;27(3):321-8.
26. Abelha FJ, Botelho M, Fernandes V, Barros H. Quality of life and mortality assessment in patients with major cardiac events in the postoperative period. *Revista brasileira de anestesiologia*. 2010;60(3):268-84.
27. Lobo SM, Rezende E, Knibel MF, Silva NB, Paramo JA, Nacul FE, et al. Early determinants of death due to multiple organ failure after noncardiac surgery in high-risk patients. *Anesthesia and analgesia*. 2011;112(4):877-83.
28. Abelha FJ, Luis C, Veiga D, Parente D, Fernandes V, Santos P, et al. Outcome and quality of life in patients with postoperative delirium during an ICU stay following major surgery. *Critical care*. 2013;17(5):R257.
29. Abelha FJ, Botelho M, Fernandes V, Barros H. Determinants of postoperative acute kidney injury. *Critical care*. 2009;13(3):R79.
30. Rhodes A, Moreno RP, Metnitz B, Hochrieser H, Bauer P, Metnitz P. Epidemiology and outcome following post-surgical admission to critical care. *Intensive care medicine*. 2011;37(9):1466-72.
31. Sakr Y, Vincent JL, Ruokonen E, Pizzamiglio M, Installé E, Reinhart K, et al. Sepsis and organ system failure are major determinants of post-intensive care unit mortality. *Journal of critical care*. 2008;23(4):475-83.
32. Xara D, Santos A, Abelha F. Adverse Respiratory Events in a Post-Anesthesia Care Unit. *Archivos de bronconeumologia*. 2015;51(2):69-75.
33. Elias AC, Matsuo T, Grion CM, Cardoso LT, Verri PH. Incidence and risk factors for sepsis in surgical patients: a cohort study. *Journal of critical care*. 2012;27(2):159-66.
34. Vascular Events In Noncardiac Surgery Patients Cohort Evaluation Study I, Devereaux PJ, Chan MT, Alonso-Coello P, Walsh M, Berwanger O, et al. Association between postoperative troponin levels and 30-day mortality among patients undergoing noncardiac surgery. *Jama*. 2012;307(21):2295-304.

35. Weissman C, Klein N. The importance of differentiating between elective and emergency postoperative critical care patients. *Journal of critical care*. 2008;23(3):308-16.
36. Choi WI, Shehu E, Lim SY, Koh SO, Jeon K, Na S, et al. Markers of poor outcome in patients with acute hypoxemic respiratory failure. *Journal of critical care*. 2014;29(5):797-802.
37. Cooke CR, Kahn JM, Caldwell E, Okamoto VN, Heckbert SR, Hudson LD, et al. Predictors of hospital mortality in a population-based cohort of patients with acute lung injury. *Critical care medicine*. 2008;36(5):1412-20.
38. de Jonge E, Peelen L, Keijzers PJ, Joore H, de Lange D, van der Voort PH, et al. Association between administered oxygen, arterial partial oxygen pressure and mortality in mechanically ventilated intensive care unit patients. *Critical care*. 2008;12(6):R156.
39. Chahoud J, Semaan A, Almoosa KF. Ventilator-associated events prevention, learning lessons from the past: A systematic review. *Heart & lung : the journal of critical care*. 2015.
40. Kallet RH, Matthay MA. Hyperoxic acute lung injury. *Respiratory care*. 2013;58(1):123-41.
41. Darmon M, Diconne E, Souweine B, Ruckly S, Adrie C, Azoulay E, et al. Prognostic consequences of borderline dysnatremia: pay attention to minimal serum sodium change. *Critical care*. 2013;17(1):R12.
42. Funk GC, Lindner G, Druml W, Metnitz B, Schwarz C, Bauer P, et al. Incidence and prognosis of dysnatremias present on ICU admission. *Intensive care medicine*. 2010;36(2):304-11.
43. Darmon M, Timsit JF, Francais A, Nguile-Makao M, Adrie C, Cohen Y, et al. Association between hypernatraemia acquired in the ICU and mortality: a cohort study. *Nephrology, dialysis, transplantation : official publication of the European Dialysis and Transplant Association - European Renal Association*. 2010;25(8):2510-5.
44. Waite MD, Fuhrman SA, Badawi O, Zuckerman IH, Franey CS. Intensive care unit-acquired hypernatremia is an independent predictor of increased mortality and length of stay. *Journal of critical care*. 2013;28(4):405-12.
45. Stelfox HT, Ahmed SB, Khandwala F, Zygun D, Shahpori R, Laupland K. The epidemiology of intensive care unit-acquired hyponatraemia and hypernatraemia in medical-surgical intensive care units. *Critical care*. 2008;12(6):R162.

46. Lindner G, Funk GC. Hyponatremia in critically ill patients. *Journal of critical care*. 2013;28(2):216 e11-20.
47. Martin MJ, FitzSullivan E, Salim A, Berne TV, Towfigh S. Use of serum bicarbonate measurement in place of arterial base deficit in the surgical intensive care unit. *Archives of surgery*. 2005;140(8):745-51.
48. Gunnerson KJ, Saul M, He S, Kellum JA. Lactate versus non-lactate metabolic acidosis: a retrospective outcome evaluation of critically ill patients. *Critical care*. 2006;10(1):R22.
49. Surbatovic M, Radakovic S, Jevtic M, Filipovic N, Romic P, Popovic N, et al. Predictive value of serum bicarbonate, arterial base deficit/excess and SAPS III score in critically ill patients. *General physiology and biophysics*. 2009;28 Spec No:271-6.
50. Meregalli A, Oliveira RP, Friedman G. Occult hypoperfusion is associated with increased mortality in hemodynamically stable, high-risk, surgical patients. *Critical care*. 2004;8(2):R60-5.
51. Liborio AB, Noritomi DT, Leite TT, de Melo Bezerra CT, de Faria ER, Kellum JA. Increased serum bicarbonate in critically ill patients: a retrospective analysis. *Intensive care medicine*. 2015;41(3):479-86.
52. Paz Y, Zegerman A, Sorkine P, Matot I. Severe acidosis does not predict fatal outcomes in intensive care unit patients: a retrospective analysis. *Journal of critical care*. 2014;29(2):210-3.
53. Naughton C, Feneck RO. The impact of age on 6-month survival in patients with cardiovascular risk factors undergoing elective non-cardiac surgery. *International journal of clinical practice*. 2007;61(5):768-76.
54. Haynes AB, Weiser TG, Berry WR, Lipsitz SR, Breizat AH, Dellinger EP, et al. A surgical safety checklist to reduce morbidity and mortality in a global population. *The New England journal of medicine*. 2009;360(5):491-9.
55. Thompson JS, Baxter BT, Allison JG, Johnson FE, Lee KK, Park WY. Temporal patterns of postoperative complications. *Archives of surgery*. 2003;138(6):596-602.
56. Cuthbertson BH, Campbell MK, Stott SA, Vale L, Norrie J, Kinsella J, et al. A pragmatic multi-centre randomised controlled trial of fluid loading and level of dependency in high-risk surgical patients undergoing major elective surgery: trial protocol. *Trials*. 2010;11:41.

57. Wolters U, Wolf T, Stutzer H, Schroder T. ASA classification and perioperative variables as predictors of postoperative outcome. *British journal of anaesthesia*. 1996;77(2):217-22.
58. Moonesinghe SR, Mythen MG, Das P, Rowan KM, Grocott MP. Risk stratification tools for predicting morbidity and mortality in adult patients undergoing major surgery: qualitative systematic review. *Anesthesiology*. 2013;119(4):959-81.
59. Basques BA, Fu MC, Buerba RA, Bohl DD, Golinvaux NS, Grauer JN. Using the ACS-NSQIP to identify factors affecting hospital length of stay after elective posterior lumbar fusion. *Spine*. 2014;39(6):497-502.

Tables

Table 1: Univariate analysis of mortality predictors in SICU – patients' characteristics

Variables	Total N=4398	Survival group N=4338	Mortality group N=60	p value
Gender, n (%)				
Male	2681 (61.0)	2642 (60.9)	39 (65.0)	0.518 ^{a)}
Female	1717 (39.0)	1696 (39.1)	21 (35.0)	
Age, median (IQR)	65.0 (54.0-74.0)	65.0 (53.0-74.0)	72.5 (59.5-79.8)	<0.001 ^{b)}
Type of admission, n (%)				
Elective surgery	3827 (87.0)	3803 (87.7)	24 (40.0)	<0.001 ^{a)}
Non-elective surgery	571 (13.0)	535 (12.3)	36 (60.0)	
Mechanical ventilation on admission, n (%)	1341 (30.5)	1291 (29.8)	50 (83.3)	<0.001 ^{a)}
Renal insufficiency, n (%)	285 (6.5)	264 (6.1)	21 (35.0)	<0.001 ^{f)}
Organ insufficiency, n (%)	682 (15.5)	658 (15.2)	24 (40.0)	<0.001 ^{a)}
Hematocrit, median (IQR)	33.0 (29.8-36.3)	33.0 (29.9-36.4)	28.8 (22.9-33.0)	<0.001 ^{b)}
Body temperature, median (IQR)	35.4 (34.6-36.0)	35.8 (34.6-36.0)	34.0 (33.0-35.3)	<0.001 ^{b)}
Systolic pressure, median (IQR)	122.0 (102.0-144.0)	122.0 (102.0-144.0)	76.5 (66.0-88.8)	<0.001 ^{b)}
Mean arterial pressure, median (IQR)	85.0 (71.0-96.0)	85.0 (71.0-96.0)	53.0 (47.3-63.0)	<0.001 ^{b)}
Heart rate, median (IQR)	83 (69-98)	83 (68-98)	112 (88-133)	<0.001 ^{b)}
Respiratory rate, median (IQR)	14 (12-16)	14 (12-16)	16 (14-16)	<0.001 ^{b)}
Serum urea, median (IQR)	30.0 (20.0-40.0)	30.0 (20.0-40.0)	45.0 (23.5-70.0)	0.001 ^{b)}
Serum creatinine, median (IQR)	8.3 (6.5-11.0)	8.2 (6.5-11.0)	15.6 (9.0-25.3)	<0.001 ^{b)}
Total bilirubin, median (IQR)	4.0 (1.0-7.0)	4.0 (1.0-7.0)	6.0 (4.0-10.8)	<0.001 ^{b)}
FiO2, median (IQR)	0.40 (0.35-0.40)	0.40 (0.34-0.40)	0.52 (0.40-1.00)	<0.001 ^{b)}
PaO2, median (IQR)	100.0 (100.0-110.0)	100.0 (100.0-110.0)	98.0 (75.5-138.6)	0.039 ^{b)}
PaCO2, median (IQR)	39.5 (35.0-45.0)	39.4 (35.0-45.0)	42.7 (36.0-54.0)	0.001 ^{b)}
Serum bicarbonate, median (IQR)	22.0 (21.0-24.0)	22.0 (21.0-24.0)	19.4 (17.0-22.0)	<0.001 ^{b)}
pH, median (IQR)	7.40 (7.35-7.40)	7.40 (7.35-7.40)	7.28 (7.17-7.35)	<0.001 ^{b)}
Serum potassium, median (IQR)	3.80 (3.40-4.10)	3.80 (3.40-4.10)	3.90 (3.13-4.45)	0.806 ^{b)}
Serum sodium, median (IQR)	140 (137-142)	140 (137-142)	145 (140-152)	<0.001 ^{b)}
Leucocytes count, median (IQR)	11.0 (8.0-14.0)	11.0 (8.0-11.0)	9.5 (4.0-19.0)	0.230 ^{b)}

Glasgow coma scale (<9), n (%)	54 (1.2)	46 (1.1)	8 (13.3)	<0.001 ^{f)}
--------------------------------	----------	----------	----------	----------------------

APACHE II: Acute Physiology and Chronic Health Evaluation; IQR: Interquartile range (P25-P75);

SAPS II: Simplified Acute Physiology Score; SICU: Surgical Intensive Care Unit

^{a)} Chi-Square test

^{b)} Mann-Whitney test

^{f)} p value with Fisher's Exact Test

Table 2: Univariate analysis of mortality predictors in SICU - severity of disease scores, length of stay and major cardiac event.

Variables	Total N=4398	Survival group N=4338	Mortality group N=60	p value
APACHE II, median (IQR)	8.0 (6.0-12.0)	8.0 (6.0-12.0)	22.0 (19.0-26.0)	<0.001 ^{b)}
SAPS II, median (IQR)	18.0 (13.3-26.7)	18.0 (13.3-25.0)	43.7 (37.8-57.8)	<0.001 ^{b)}
SICU LOS(hours), median (IQR)	20.0 (16.0-42.0)	20.0 (16.0-41.0)	46.0 (19.5-82.8)	<0.001 ^{b)}
Major cardiac event , n (%)	107 (2.4)	90 (2.1)	17 (28.3)	<0.001 ^{f)}

APACHE II: Acute Physiology and Chronic Health Evaluation; IQR: Interquartile range (P25-P75); LOS: length of stay; SAPS II: Simplified Acute Physiology Score; SICU: Surgical Intensive Care Unit

^{b)} Mann-Whitney test

^{f)} p value with Fisher's Exact Test

Table 3: Univariate analysis of mortality predictors in SICU - Criteria developed by Lee et al

Variables	Total N=4398	Survival group N=4338	Mortality group N=60	p value
High-risk surgery, n (%)	2382 (54.2)	2334 (53.8)	48 (80.0)	<0.001 ^{a)}
History of ischemic heart disease, n (%)	617 (14.0)	607 (14.0)	10 (16.7)	0.554 ^{a)}
History of congestive heart disease, n (%)	691 (15.7)	673 (15.5)	18 (30.0)	0.002 ^{a)}
Preoperative insulin therapy, n (%)	215 (4.9)	213 (4.9)	2 (3.3)	1 ^{f)}
Preoperative serum creatinine >2.0 mg/dL, n (%)	281 (6.4)	272 (6.3)	9 (15.0)	0.013 ^{f)}
History of cerebrovascular disease, n (%)	559 (12.7)	548 (12.6)	11 (18.3)	0.188 ^{a)}
RCRI ≥ 2 , n (%)	328 (7.5)	317 (7.3)	11 (18.3)	0.004 ^{f)}

RCRI: Revised Cardiac Risk Index; SICU: Surgical Intensive Care Unit

^{a)} Chi-Square test

^{f)} p value with Fisher's Exact Test

Table 4: Multiple logistic regression analysis for independent predictors of mortality in SICU

Variables	OR ^{a)} (95% IC)	p ^{b)} value
APACHE II	1.25 (1.18- 1.31)	<0.001
Type of admission	3.88 (2.02- 7.46)	<0.001
FIO2	21.25 (4.75- 95.16)	<0.001
Serum bicarbonate	0.88 (0.82- 0.95)	0.001
Serum sodium	1.06 (1.01- 1.11)	0.01

APACHE II: Acute Physiology and Chronic Health Evaluation; CI: confidence interval; OR: odds ratio; SICU: Surgical Intensive Care Unit;

^{a)} Adjusted for age, mechanical ventilation on admission, renal insufficiency, organ insufficiency, hematocrit, body temperature, systolic pressure, mean arterial pressure, heart rate, respiratory rate, serum urea, serum creatinine, total bilirubin, FiO2, PaCO2, serum bicarbonate, pH, serum sodium, glasgow coma scale and SAPS II

^{b)} A logistic regression analysis with inclusion severity of disease scoring systems' variables with $p \leq 0.001$ was used.

Com toda a sinceridade agradeço ao meu orientador, Prof. Doutor Fernando Abelha, pela oportunidade de realizar este trabalho e por todo o apoio e aconselhamento durante o seu desenvolvimento. Agradeço ainda toda a compreensão e disponibilidade. Obrigada!



JOURNAL OF CRITICAL CARE

Improving Patient Care by Integrating Critical Care Systems Knowledge into Practice Behavior

AUTHOR INFORMATION PACK

TABLE OF CONTENTS

ISSN: 0883-9441

•	Description	p.1
•	Impact Factor	p.1
•	Editorial Board	p.1
•	Guide for Authors	p.3

DESCRIPTION

The *Journal of Critical Care*, the official publication of the World Federation of Societies of Intensive and Critical Care Medicine (WFSICCM), is a leading international, peer-reviewed journal providing original research, review articles, tutorials, and invited articles for physicians and allied health professionals involved in treating the **critically ill**. The Journal aims to improve patient care by furthering understanding of **health systems** research and its integration into **clinical practice**.

The Journal will include articles which discuss:

All aspects of **health services** research in critical care System based practice in **anesthesiology**, **perioperative** and **critical care medicine** The interface between anesthesiology, critical care medicine and pain Integrating **intraoperative management** in preparation for postoperative critical care management and recovery Optimizing patient management, i.e., exploring the interface between evidence-based principles or clinical insight into management and care of complex patients The team approach in the OR and ICU System-based research **Medical ethics Technology** in medicine Seminars discussing current, state of the art, and sometimes controversial topics in anesthesiology, critical care medicine, and professional education Residency Education: Providing a series of clinically relevant tutorials from experienced practitioners focusing on the six core competencies: Medical Knowledge Professionalism Patient Care Interpersonal and communication skills Case based learning and improvement Systems based practice

The [editorial board](#) represents an international cross section of individuals actively involved in the disciplines of Critical Care Medicine and Anesthesiology.

IMPACT FACTOR

2013: 2.191 © Thomson Reuters Journal Citation Reports 2014

EDITORIAL BOARD

Editor-in-Chief:

Philip D. Lumb, MB, BS, MCCM, USA

Editorial Board

André Amaral MD, Brazil

Jan Bakker MD, PhD, The Netherlands

Satish Bhagwanjee, USA

Timothy George Buchman MD, PhD, USA

Jean Carlet MD, France

Jack Chen MBBS, PhD, Australia

Sanjeev Chhangani, USA
Jason Christie MD, USA
Gilles Clermont MD, USA
Edmond Cohen, USA
Deborah Cook MD, Canada
Heidi Frankel, USA
Elizabeth Frost MD, USA
Cristina Granja MD, PhD, Portugal
Antonino Gullo MD, Italy
Steven Haddy, USA
Graeme K. Hart MD, Australia
Kenneth Mark Hillman MBBS, Australia
Ramona O. Hopkins PhD, USA
Catherine Hough MD, MSc, USA
Michael Imhoff MD, Germany
Dimitrios Karakitsos, Greece
D.J. (Jim) Kutsogiannis MD, MHS, Canada
Christoph Lehmann MD, USA
Brenda Lynn Morgan RN, BScN, MSc, Canada
Victor Moroz MD, Russia
Edmund Neugebauer PhD, Germany
Stephen M. Pastores, USA
Rajesh Patel MD, MEd, USA
Michael Pinsky MD, USA
Adrienne G. Randolph MD, MSc, USA
Kathryn Rowan PhD, United Kingdom
Damon Scales MD, Canada
Andrew J.E. Seely MD, Canada
Ram Shanmugam PhD, USA
Kevin N. Sheth, USA
Khalid Shukri, Saudi Arabia
Charles L. Sprung MD, Israel
Jean-Louis Vincent MD, PhD, Belgium
Yoram Vodovotz PhD, USA
Vladimir Zelman MD, PhD, USA

Managing Editor:

Diane McIntee MS

World Federation Council Members (WFSICCM)

Jean-Louis Vincent, President, Belgium
John Marshall, Secretary General, Canada
Janice Zimmerman, Treasurer, United States
Edgar Jimenez, Past President, United States
Pravin Amin, India
Djillai Annane, France
Llius Blanch, Spain
Guillermo Castorena, Mexico
Raffaele de Gaudio, Italy
Bin Du, China
Younsuck Koh, Korea
John Myburgh, Australia
Masaji Nishimura, Japan
Alvaro Rea-Neto, Brazil
Arzu Topeli, Turkey
Sebastian Ugarte, Chile

GUIDE FOR AUTHORS

The Journal of Critical Care provides a forum for the publication of original peer-reviewed articles with the goal of improving patient care by integrating critical care systems knowledge into practice behavior. The journal represents the World Federation of Societies of Intensive and Critical Care Medicine (WFSICCM), an organization of 42 national intensive/critical care societies representing some 32,000 physicians and allied health professionals. With this responsibility to the WFSICCM comes an international focus in systems research in constrained resource environments.

We accept research articles and review articles as well as those in a seminar or tutorial format. Topics covered are all aspects of Health Services Research, the interface of critical care, anesthesiology, and pain, as well as tutorials for residency education core competencies.

For the seminar format, the articles should be directed to the resident or practicing healthcare professional. We are particularly interested in your up-to-date evaluation of the topic. Dealing with subjects of current and sometimes controversial educational and research themes is acceptable. The seminar format lends itself to a more informal presentation. Express your own viewpoints, but feel free to discuss other viewpoints as well. If you are uncertain, please indicate the degree of your uncertainty. We are looking for an absolutely honest evaluation of the topic.

Manuscripts are accepted for consideration on the condition that they are contributed solely to the Journal of Critical Care. No substantial part of a paper may have been or may be published elsewhere, except for an abstract of 200 words or less.

Manuscripts will be critically reviewed by the Editor with appropriate independent referees drawn from the Editorial Board and other experts. Acknowledgments to other investigators for advice or data must be substantiated by written authorization specifically granting permission to authors. Upon submission, authors will be required to fill out an ethics in publishing form. All submissions must also include a conflict of interest statement.

BEFORE YOU BEGIN

Plagiarism

Plagiarism and duplicate submission are serious acts of misconduct. Plagiarism is defined as unreferenced use of published or unauthorized use of unpublished ideas, and may occur at any stage of planning, researching, writing, or publication. Plagiarism takes many forms, from 'passing off' another's paper as the author's own paper, to copying or paraphrasing substantial parts of another's paper (without attribution), to claiming results from research conducted by others. Plagiarism in all its forms constitutes unethical publishing behavior and is unacceptable. Duplicate submission/publication occurs when two or more papers, without full cross-reference, share the same hypothesis, data, discussion points, or conclusions.

Ethics in publishing

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Conflict of interest

All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. See also <http://www.elsevier.com/conflictsofinterest>. Further information and an example of a Conflict of Interest form can be found at: http://help.elsevier.com/app/answers/detail/a_id/286/p/7923.

Submission declaration

Submission of an article implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere including electronically in the same form, in English or in any other language, without the written consent of the copyright-holder.

Copyright

This journal offers authors a choice in publishing their research: Open access and Subscription.

For subscription articles

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (for more information on this and copyright, see <http://www.elsevier.com/copyright>). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. Permission of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations (please consult <http://www.elsevier.com/permissions>). If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has preprinted forms for use by authors in these cases: please consult <http://www.elsevier.com/permissions>.

For open access articles

Upon acceptance of an article, authors will be asked to complete an 'Exclusive License Agreement' (for more information see <http://www.elsevier.com/OAauthoragreement>). Permitted reuse of open access articles is determined by the author's choice of user license (see <http://www.elsevier.com/openaccesslicenses>).

Retained author rights

As an author you (or your employer or institution) retain certain rights. For more information on author rights for:

Subscription articles please see <http://www.elsevier.com/journal-authors/author-rights-and-responsibilities>.
Open access articles please see <http://www.elsevier.com/OAauthoragreement>.

Role of the funding source

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated.

Funding body agreements and policies

Elsevier has established agreements and developed policies to allow authors whose articles appear in journals published by Elsevier, to comply with potential manuscript archiving requirements as specified as conditions of their grant awards. To learn more about existing agreements and policies please visit <http://www.elsevier.com/fundingbodies>.

Open access

This journal offers authors a choice in publishing their research:

Open access

- Articles are freely available to both subscribers and the wider public with permitted reuse
- An open access publication fee is payable by authors or their research funder

Subscription

- Articles are made available to subscribers as well as developing countries and patient groups through our access programs (<http://www.elsevier.com/access>)
- No open access publication fee

All articles published open access will be immediately and permanently free for everyone to read and download. Permitted reuse is defined by your choice of one of the following Creative Commons user licenses:

Creative Commons Attribution-NonCommercial-ShareAlike (CC BY-NC-SA): for non-commercial purposes, lets others distribute and copy the article, to create extracts, abstracts and other revised versions, adaptations or derivative works of or from an article (such as a translation), to include in a collective work (such as an anthology), to text and data mine the article, as long as they credit the author(s), do not represent the author as endorsing their adaptation of the article, do not modify the article in such a way as to damage the author's honor or reputation, and license their new adaptations or creations under identical terms (CC BY-NC-SA).

Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND): for non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.

Elsevier has established agreements with funding bodies, <http://www.elsevier.com/fundingbodies>. This ensures authors can comply with funding body open access requirements, including specific user licenses, such as CC BY. Some authors may also be reimbursed for associated publication fees. If you need to comply with your funding body policy, you can apply for the CC BY license after your manuscript is accepted for publication.

To provide open access, this journal has a publication fee which needs to be met by the authors or their research funders for each article published open access.

Your publication choice will have no effect on the peer review process or acceptance of submitted articles.

The open access publication fee for this journal is **\$2,500**, excluding taxes. Learn more about Elsevier's pricing policy: <http://www.elsevier.com/openaccesspricing>.

Language (usage and editing services)

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the English Language Editing service available from Elsevier's WebShop (<http://webshop.elsevier.com/languageediting/>) or visit our customer support site (<http://support.elsevier.com>) for more information.

Submission

Submission to this journal proceeds totally online and you will be guided stepwise through the creation and uploading of your files. The system automatically converts source files to a single PDF file of the article, which is used in the peer-review process. Please note that even though manuscript source files are converted to PDF files at submission for the review process, these source files are needed for further processing after acceptance. All correspondence, including notification of the Editor's decision and requests for revision, takes place by e-mail removing the need for a paper trail.

Submit your article

Please submit your article via <http://ees.elsevier.com/jcrc/>.

NIH Public Access Policy Compliance

To comply with the NIH Public Access Policy, Elsevier will deposit to PubMed Central (PMC) author manuscripts on behalf of authors reporting NIH funded research. The NIH policy requires that NIH-funded authors submit to PubMed Central (PMC), or have submitted on their behalf, their peer-reviewed author manuscripts, to appear on PMC no later than 12 months after final publication. Elsevier will send to PMC the final peer-reviewed manuscript, which was accepted for publication and sent to Elsevier's production department, and that reflects any author-agreed changes made in response to peer-review comments. Elsevier will authorize the author manuscript's public access posting 12 months after final publication. Following the deposit by Elsevier, authors will receive further communications from the NIH with respect to the submission. Note: Authors must declare their NIH funding (or the other funding bodies listed below) when completing the copyright transfer form.

PREPARATION

Use of word processing software

It is important that the file be saved in the native format of the word processor used. The text should be in single-column format. Keep the layout of the text as simple as possible. Most formatting codes will be removed and replaced on processing the article. In particular, do not use the word processor's options to justify text or to hyphenate words. However, do use bold face, italics, subscripts, superscripts etc. When preparing tables, if you are using a table grid, use only one grid for each individual table and not a grid for each row. If no grid is used, use tabs, not spaces, to align columns. The electronic text should be prepared in a way very similar to that of conventional manuscripts (see also the Guide to Publishing with Elsevier: <http://www.elsevier.com/guidepublication>). Note that source files of figures, tables and text graphics will be required whether or not you embed your figures in the text. See also the section on Electronic artwork.

To avoid unnecessary errors you are strongly advised to use the 'spell-check' and 'grammar-check' functions of your word processor.

Embedded math equations

If you are submitting an article prepared with Microsoft Word containing embedded math equations then please read this related support information (http://support.elsevier.com/app/answers/detail/a_id/302/).

Article structure

Subdivision - unnumbered sections

Divide your article into clearly defined sections. Each subsection is given a brief heading. Each heading should appear on its own separate line. Subsections should be used as much as possible when cross-referencing text: refer to the subsection by heading as opposed to simply 'the text'.

Introduction

State the objectives of the work and provide an adequate background, avoiding a detailed literature survey or a summary of the results.

Material and methods

Provide sufficient detail to allow the work to be reproduced. Methods already published should be indicated by a reference: only relevant modifications should be described.

Results

Results should be clear and concise.

Discussion

This should explore the significance of the results of the work, not repeat them. A combined Results and Discussion section is often appropriate. Avoid extensive citations and discussion of published literature.

Conclusions

The main conclusions of the study may be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.

Essential title page information

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Where the family name may be ambiguous (e.g., a double name), please indicate this clearly. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. **Ensure that phone numbers (with country and area code) are provided in addition to the e-mail address and the complete postal address. Contact details must be kept up to date by the corresponding author.**
- **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

Abstract

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

Abstract

Note: Research papers should contain a structured abstract of no more than 200 words which contain the following:

1. Purpose: Why was this study done?
2. Materials and Methods: What was the source of the data generated? How was it obtained?
3. Results: Findings should be objectively reported and statistical significance indicated (if appropriate).

4. Conclusions: Do not include a summary at the end of the paper.

Graphical abstract

Although a graphical abstract is optional, its use is encouraged as it draws more attention to the online article. The graphical abstract should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of 531 × 1328 pixels (h × w) or proportionally more. The image should be readable at a size of 5 × 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. See <http://www.elsevier.com/graphicalabstracts> for examples.

Authors can make use of Elsevier's Illustration and Enhancement service to ensure the best presentation of their images and in accordance with all technical requirements: [Illustration Service](#).

Keywords

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

Abbreviations

Define abbreviations that are not standard in this field in a footnote to be placed on the first page of the article. Such abbreviations that are unavoidable in the abstract must be defined at their first mention there, as well as in the footnote. Ensure consistency of abbreviations throughout the article.

Acknowledgements

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

Embedded math equations

If you are submitting an article prepared with Microsoft Word containing embedded math equations then please read this related support information (http://support.elsevier.com/app/answers/detail/a_id/302/).

Artwork

Electronic artwork

General points

- Make sure you use uniform lettering and sizing of your original artwork.
- Embed the used fonts if the application provides that option.
- Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Size the illustrations close to the desired dimensions of the printed version.
- Submit each illustration as a separate file.

A detailed guide on electronic artwork is available on our website:

<http://www.elsevier.com/artworkinstructions>

You are urged to visit this site; some excerpts from the detailed information are given here.

Formats

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.

Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings, embed all used fonts.

TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.

TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.

TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

Please do not:

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

Color artwork

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color on the Web (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or on the Web only. For further information on the preparation of electronic artwork, please see <http://www.elsevier.com/artworkinstructions>.

Please note: Because of technical complications that can arise by converting color figures to 'gray scale' (for the printed version should you not opt for color in print) please submit in addition usable black and white versions of all the color illustrations.

Illustration services

Elsevier's WebShop (<http://webshop.elsevier.com/illustrationservices>) offers Illustration Services to authors preparing to submit a manuscript but concerned about the quality of the images accompanying their article. Elsevier's expert illustrators can produce scientific, technical and medical-style images, as well as a full range of charts, tables and graphs. Image 'polishing' is also available, where our illustrators take your image(s) and improve them to a professional standard. Please visit the website to find out more.

Figure captions

Ensure that each illustration has a caption. Supply captions separately, not attached to the figure. A caption should comprise a brief title (**not** on the figure itself) and a description of the illustration. Keep text in the illustrations themselves to a minimum but explain all symbols and abbreviations used.

Tables

Number tables consecutively in accordance with their appearance in the text. Place footnotes to tables below the table body and indicate them with superscript lowercase letters. Avoid vertical rules. Be sparing in the use of tables and ensure that the data presented in tables do not duplicate results described elsewhere in the article.

References

Citation in text

Please ensure that every reference cited in the text is also present in the reference list (and vice versa). Any references cited in the abstract must be given in full. Unpublished results and personal communications are not recommended in the reference list, but may be mentioned in the text. If these references are included in the reference list they should follow the standard reference style of the journal and should include a substitution of the publication date with either 'Unpublished results' or 'Personal communication'. Citation of a reference as 'in press' implies that the item has been accepted for publication.

Reference links

Increased discoverability of research and high quality peer review are ensured by online links to the sources cited. In order to allow us to create links to abstracting and indexing services, such as Scopus, CrossRef and PubMed, please ensure that data provided in the references are correct. Please note that incorrect surnames, journal/book titles, publication year and pagination may prevent link creation. When copying references, please be careful as they may already contain errors. Use of the DOI is encouraged.

Web references

As a minimum, the full URL should be given and the date when the reference was last accessed. Any further information, if known (DOI, author names, dates, reference to a source publication, etc.), should also be given. Web references can be listed separately (e.g., after the reference list) under a different heading if desired, or can be included in the reference list.

References in a special issue

Please ensure that the words 'this issue' are added to any references in the list (and any citations in the text) to other articles in the same Special Issue.

Reference style

Text: Indicate references by number(s) in square brackets in line with the text. The actual authors can be referred to, but the reference number(s) must always be given.

List: Number the references (numbers in square brackets) in the list in the order in which they appear in the text.

Examples:

Reference to a journal publication:

[1] Van der Geer J, Hanraads JAJ, Lupton RA. The art of writing a scientific article. *J Sci Commun* 2010;163:51–9.

Reference to a book:

[2] Strunk Jr W, White EB. *The elements of style*. 4th ed. New York: Longman; 2000.

Reference to a chapter in an edited book:

[3] Mettam GR, Adams LB. How to prepare an electronic version of your article. In: Jones BS, Smith RZ, editors. *Introduction to the electronic age*, New York: E-Publishing Inc; 2009, p. 281–304.

Note shortened form for last page number. e.g., 51–9, and that for more than 6 authors the first 6 should be listed followed by 'et al.' For further details you are referred to 'Uniform Requirements for Manuscripts submitted to Biomedical Journals' (*J Am Med Assoc* 1997;277:927–34) (see also http://www.nlm.nih.gov/bsd/uniform_requirements.html).

Journal abbreviations source

Journal names should be abbreviated according to the List of Title Word Abbreviations: <http://www.issn.org/services/online-services/access-to-the-ltwa/>.

Video data

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. In order to ensure that your video or animation material is directly usable, please provide the files in one of our recommended file formats with a preferred maximum size of 50 MB. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including ScienceDirect: <http://www.sciencedirect.com>. Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our video instruction pages at <http://www.elsevier.com/artworkinstructions>. Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

AudioSlides

The journal encourages authors to create an AudioSlides presentation with their published article. AudioSlides are brief, webinar-style presentations that are shown next to the online article on ScienceDirect. This gives authors the opportunity to summarize their research in their own words and to help readers understand what the paper is about. More information and examples are available at <http://www.elsevier.com/audioslides>. Authors of this journal will automatically receive an invitation e-mail to create an AudioSlides presentation after acceptance of their paper.

Supplementary data

Elsevier accepts electronic supplementary material to support and enhance your scientific research. Supplementary files offer the author additional possibilities to publish supporting applications, high-resolution images, background datasets, sound clips and more. Supplementary files supplied will be published online alongside the electronic version of your article in Elsevier Web products, including ScienceDirect: <http://www.sciencedirect.com>. In order to ensure that your submitted material is directly usable, please provide the data in one of our recommended file formats. Authors should submit the material in electronic format together with the article and supply a concise and descriptive caption for each file. For more detailed instructions please visit our artwork instruction pages at <http://www.elsevier.com/artworkinstructions>.

Submission checklist

The following list will be useful during the final checking of an article prior to sending it to the journal for review. Please consult this Guide for Authors for further details of any item.

Ensure that the following items are present:

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address
- Phone numbers

All necessary files have been uploaded, and contain:

- Keywords
- All figure captions
- All tables (including title, description, footnotes)

Further considerations

- Manuscript has been 'spell-checked' and 'grammar-checked'
- References are in the correct format for this journal
- All references mentioned in the Reference list are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources (including the Web)
- Color figures are clearly marked as being intended for color reproduction on the Web (free of charge) and in print, or to be reproduced in color on the Web (free of charge) and in black-and-white in print
- If only color on the Web is required, black-and-white versions of the figures are also supplied for printing purposes

For any further information please visit our customer support site at <http://support.elsevier.com>.

AFTER ACCEPTANCE***Use of the Digital Object Identifier***

The Digital Object Identifier (DOI) may be used to cite and link to electronic documents. The DOI consists of a unique alpha-numeric character string which is assigned to a document by the publisher upon the initial electronic publication. The assigned DOI never changes. Therefore, it is an ideal medium for citing a document, particularly 'Articles in press' because they have not yet received their full bibliographic information. Example of a correctly given DOI (in URL format; here an article in the journal *Physics Letters B*):

<http://dx.doi.org/10.1016/j.physletb.2010.09.059>

When you use a DOI to create links to documents on the web, the DOIs are guaranteed never to change.

Proofs

One set of page proofs (as PDF files) will be sent by e-mail to the corresponding author (if we do not have an e-mail address then paper proofs will be sent by post) or, a link will be provided in the e-mail so that authors can download the files themselves. Elsevier now provides authors with PDF proofs which can be annotated; for this you will need to download Adobe Reader version 9 (or higher) available free from <http://get.adobe.com/reader>. Instructions on how to annotate PDF files will accompany the proofs (also given online). The exact system requirements are given at the Adobe site: <http://www.adobe.com/products/reader/tech-specs.html>.

If you do not wish to use the PDF annotations function, you may list the corrections (including replies to the Query Form) and return them to Elsevier in an e-mail. Please list your corrections quoting line number. If, for any reason, this is not possible, then mark the corrections and any other comments (including replies to the Query Form) on a printout of your proof and return by fax, or scan the pages and e-mail, or by post. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. We will do everything possible to get your article published quickly and accurately. It is important to ensure that all corrections are sent back to us in one communication: please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

Other Funding Body Policies

Elsevier has also worked with the following funding bodies to ensure that our authors can comply with their policies:

- Arthritis Research Campaign (UK)
- British Heart Foundation (UK)
- Cancer Research (UK)
- Chief Scientist Office
- Department of Health (UK)
- Howard Hughes Medical Institute (US)

- Medical Research Council (UK)
- Wellcome Trust (UK)

For full details on how these policies are implemented, please see complete information at:
<http://www.elsevier.com/wps/find/authorsview.authors/fundingbodyagreements>.

AUTHOR INQUIRIES

You can track your submitted article at http://help.elsevier.com/app/answers/detail/a_id/89/p/8045/.
You can track your accepted article at <http://www.elsevier.com/trackarticle>. You are also welcome to contact Customer Support via <http://support.elsevier.com>.

© Copyright 2014 Elsevier | <http://www.elsevier.com>

Declaração

Para os devidos efeitos se declara que o Prof. Doutor Fernando José Pereira Alves Abelha entregou um projecto de investigação denominado “*Mortalidade e complicações cardíacas no pós-operatório numa população cirúrgica crítica*” para ser apreciado pela Comissão de Ética para a Saúde do Hospital de S. João na Reunião Plenária de 25 de Novembro de 2014, presidida pelo Prof. Doutor Filipe Almeida, e o mesmo foi aprovado com o parecer nº 248-2014 em 25 de Novembro de 2014.

Centro Hospitalar de S. João, 15 de Dezembro de 2014

O Secretário da Comissão de Ética para a Saúde


